

A New Changhee q -Euler Numbers and Polynomials Associated with p -Adic q -Integrals

SEOG-HOON RIM

Department of Mathematics Education, Kyungpook National University, Daegu 702-701, Korea

e-mail : `shrim@knu.ac.kr`

TAEKYUN KIM

EECS, Kyungpook National University, Taegu 702-701, S.Korea

e-mail : `tkim@knu.ac.kr`, `tkim64@hanmail.net`

ABSTRACT. Using non-archimedean q -integrals on $\mathbb{Z}_p$ defined in [15, 16], we define a new Changhee q -Euler polynomials and numbers which are different from those of Kim [7] and Carlitz [2]. We define generating functions of multiple q -Euler numbers and polynomials. Furthermore we construct multivariate Hurwitz type zeta function which interpolates the multivariate q -Euler numbers or polynomials at negative integers.

1. Introduction

Let p be a fixed odd prime in this paper. Throughout this paper, the symbols $\mathbb{Z}$, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$, denote the ring of rational integers, the ring of p -adic integers, the field of p -adic numbers, the complex number field, and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let $\nu_p(p)$ be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$. When one speaks of q -extension, q can be regarded as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$; it is always clear from the context. If $q \in \mathbb{C}$, then one usually assumes that $|q| < 1$. If $q \in \mathbb{C}_p$, then one usually assumes that $|q - 1|_p < p^{-\frac{1}{p-1}}$, and hence $q^x = \exp(x \log q)$ for $x \in \mathbb{Z}_p$. In this paper, we use the below notation

$$[x]_q = \frac{1 - q^x}{1 - q}, \quad (a : q)_n = (1 - a)(1 - aq) \cdots (1 - aq^{n-1}), \quad \text{cf. [3, 4, 5, 6, 7, 8].}$$

Note that $\lim_{q \rightarrow 1} [x]_q = x$ for any x with $|x|_p \leq 1$ in the p -adic case. For a fixed

Key words and phrases: 11B68, 11S40.

2000 Mathematics Subject Classification: p -adic q -integral, q -Euler numbers and polynomials.

positive integer d with $(p, d) = 1$, set

$$X = X_d = \varprojlim_N \mathbb{Z}/dp^N,$$

$$X_1 = \mathbb{Z}_p, \quad X^* = \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} a + dp\mathbb{Z}_p,$$

$$a + dp^N\mathbb{Z}_p = \{x \in X \mid x \equiv a \pmod{p^N}\},$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < dp^N$, (cf. [10,11]). We say that f is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$, and write $f \in UD(\mathbb{Z}_p)$, if the difference quotients $F_f(x, y) = \frac{f(x) - f(y)}{x - y}$ have a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$. For $f \in UD(\mathbb{Z}_p)$, let us begin with the expression

$$\frac{1}{[p^N]_q} \sum_{0 \leq j < p^N} q^j f(j) = \sum_{0 \leq j < p^N} f(j) \mu_q(j + p^N\mathbb{Z}_p), \quad \text{cf. [4, 5, 6, 8],}$$

which represents a q -analogue of Riemann sums for f . The integral of f on $\mathbb{Z}_p$ is defined as the limit of those sums (as $n \rightarrow \infty$) if this limit exists. The p -adic q -integral of a function $f \in UD(\mathbb{Z}_p)$ is defined by

$$I_q(f) = \int_X f(s) d\mu_q(x) = \int_{X_d} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[dp^N]_q} \sum_{x=0}^{dp^N-1} f(x) q^x.$$

Recently, many mathematicians studied Bernoulli and Euler numbers (see [1-24]). Using non-archimedean q -integrals on $\mathbb{Z}_p$ defined in [15, 16], we define a new Changhee q -Euler polynomials and numbers which are different from those of Kim [7] and Carlitz [2]. We define generating functions of multiple q -Euler numbers and polynomials. Furthermore we construct multivariate Hurwitz type zeta function which interpolates the multivariate q -Euler numbers or polynomials at negative integers.

2. Multivariate q -Euler numbers and polynomials

Using p -adic q -integrals on $\mathbb{Z}_p$, we now define a new q -Euler polynomials as follows:

$$\int_{\mathbb{Z}_p} q^y (x + y)^n d\mu_{-1}(y) = E_{n,q}(x).$$

Note that $\lim_{q \rightarrow 1} E_{n,q}(x) = E_n(x)$, where $E_n(x)$ are Euler polynomial which are defined by $\frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}$. In the case $x = 0$, $E_{n,q} = E_{n,q}(0)$ are called a new q -Euler numbers. And note that $\lim_{q \rightarrow 1} E_{n,q} = E_n$, where E_n are classical Euler numbers. Let $a_1, a_2, \dots, a_r, b_1, b_2, \dots, b_r$ be positive integers. Then we consider a multivariate integral as follows

$$\begin{aligned} & \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} q^{b_1 x_1 + \dots + b_r x_r} (a_1 x_1 + \dots + a_r x_r)^n d\mu_{-1}(x_1) \dots d\mu_{-1}(x_r) \\ &= E_{n,q}^{(r)}(x | a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r). \end{aligned} \quad (1)$$

Here $E_{n,q}^{(r)}(x|a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r)$ are called multivariate q -Euler polynomial of order r .

In the case $x = 0$ in (1), $E_{n,q}(0|a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r) = E_{n,q}(a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r)$ will be called multivariate q -Euler numbers of order r .

From (1), we derive the following generating function for multivariate q -Euler polynomials

$$\begin{aligned} & F_q^{(r)}(t, x|a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r) \\ &= \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x|a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r) \frac{t^n}{n!} \\ &= \frac{2^r}{(q^{b_1}e^{a_1t} + 1)(q^{b_2}e^{a_2t} + 1) \dots (q^{b_r}e^{a_rt} + 1)} e^{xt}. \end{aligned} \quad (2)$$

Note that $E_{0,q}^{(r)}(x|a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_r) = \frac{2^r}{[2]_{q^{b_1}} [2]_{q^{b_2}} \dots [2]_{q^{b_r}}}$.

Let χ be the Dirichlet character with conductor $f(=\text{odd}) \in \mathbb{N}$. Then we define the multivariate generalized q -Euler numbers attached to χ as follows:

$$\begin{aligned} & \int_X \dots \int_X \chi(x_1) \dots \chi(x_r) q^{b_1x_1 + \dots + b_rx_r} (a_1x_1 + \dots + a_rx_r)^n d\mu_{-1}(x_1) \dots d\mu_{-1}(x_r) \\ &= E_{n,\chi,q}(a_1, \dots, a_r; b_1, \dots, b_r). \end{aligned} \quad (3)$$

From (3), we can derive generating function for the multivariate generalized q -Euler numbers attached to χ ,

$$\begin{aligned} & F_{\chi,q}^{(r)}(t|a_1, \dots, a_r; b_1, \dots, b_r) \\ &= \sum_{n=0}^{\infty} E_{n,\chi,q}(a_1, \dots, a_r; b_1, \dots, b_r) \frac{t^n}{n!} \\ &= \sum_{n_1, \dots, n_r=0}^{f-1} \frac{2^r (-1)^{n_1 + \dots + n_r} q^{b_1n_1 + \dots + b_rn_r} e^{(a_1n_1 + \dots + a_rn_r)t} \chi(n_1) \dots \chi(n_r)}{(q^{fb_1}e^{fa_1t} + 1) \dots (q^{fb_r}e^{fa_rt} + 1)}, \end{aligned} \quad (4)$$

where n is odd.

Let n be odd positive integer. Form the definition of $E_{n,\chi,q}^{(r)}(a_1, \dots, a_r; b_1, \dots, b_r)$

in (3), we have the following;

$$\begin{aligned}
& E_{n,\chi,q}^{(r)}(a_1, \dots, a_r ; b_1, \dots, b_r) \\
&= \lim_{N \rightarrow \infty} \sum_{n_1, \dots, n_r=0}^{fp^N-1} \chi(n_1) \cdots \chi(n_r) (-1)^{n_1 + \dots + n_r} q^{b_1 n_1 + \dots + b_r n_r} (a_1 x_1 + \dots + a_r x_r)^n \\
&= \lim_{N \rightarrow \infty} \sum_{n_1, \dots, n_r=0}^{f-1} \sum_{x_1, \dots, x_r=0}^{p^N-1} \prod_{j=1}^r \chi(n_j + f n_j) (-1)^{\sum_{j=1}^r (n_j + f x_j)} q^{\sum_{j=1}^r b_j (n_j + f x_j)} \\
&\quad \sum_{j=1}^r a_j (n_j + f x_j)^n \\
&= f^n \sum_{n_1, \dots, n_r=0}^{f-1} (-1)^{\sum_{j=1}^r n_j} \prod_{j=1}^r \chi(n_j) q^{\sum_{j=1}^r b_j n_j} \\
&\quad \lim_{N \rightarrow \infty} \sum_{x_1, \dots, x_r=0}^{p^N-1} (-1)^{\sum_{j=1}^r (x_j)} q^{f \sum_{j=1}^r b_j x_j} \left(\frac{\sum_{j=1}^r a_j n_j}{f} + \sum_{j=1}^r a_j x_j \right)^n \\
&= f^n \sum_{n_1, \dots, n_r=0}^{f-1} (-1)^{\sum_{j=1}^r n_j} \prod_{j=1}^r \chi(n_j) q^{\sum_{j=1}^r b_j n_j} \\
&\quad \underbrace{\int_{\mathbb{Z}_r} \cdots \int_{\mathbb{Z}_p}}_{r\text{-times}} q^{f \sum_{j=1}^r b_j x_j} \left(\frac{\sum_{j=1}^n a_j x_j}{f} + \sum_{j=1}^n a_j x_j \right)^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_r) \\
&= f^n \sum_{n_1, \dots, n_r=0}^{f-1} (-1)^{\sum_{j=1}^r n_j} \prod_{j=1}^r \chi(n_j) q^{\sum_{j=1}^r b_j n_j} \\
&\quad E_{n,q^f} \left(\frac{\sum_{j=1}^n a_j x_j}{f} \mid a_1, \dots, a_r ; b_1, \dots, b_r \right).
\end{aligned}$$

Therefore we obtain;

Theorem 1. Let $a_1, \dots, a_r, b_1, \dots, b_r$ be positive integers then we have

$$\begin{aligned}
& E_{n,\chi,q}^{(r)}(a_1, \dots, a_r ; b_1, \dots, b_r) \\
&= \sum_{n_1, \dots, n_r=0}^{f-1} (-1)^{\sum_{j=1}^r n_j} \prod_{j=1}^r \chi(j) q^{\sum_{j=1}^r b_j n_j} E_{n,q^f}(a_1, \dots, a_r ; b_1, \dots, b_r), (5)
\end{aligned}$$

where f, n are odd positive integers.

3. Multivariate q -zeta functions in $\mathbb{C}$

In this section we assume $q \in \mathbb{C}$ with $|q| < 1$. Let us assume that $a_1, \dots, a_r, b_1, \dots, b_r$ are positive integers.

The purpose of this section is to study multivariate Hurwitz type zeta function which interpolates multivariate q -Euler polynomials of order r at negative integers. By (2), we easily see that

$$\begin{aligned} & F_q^{(r)}(t, x \mid a_1, \dots, a_r; b_1, \dots, b_r) \\ &= 2^r \sum_{n_1, \dots, n_r=0}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} e^{(\sum_{j=1}^r a_j n_j + x)t}. \end{aligned} \quad (6)$$

By taking derivatives of order k , on both sides of (6) we obtain the following;

Theorem 2. *Let k be positive odd integer and let $a_1, \dots, a_r, b_1, \dots, b_r$ be the positive integers. Then we have*

$$\begin{aligned} & E_{k,q}^{(r)}(x \mid a_1, \dots, a_r; b_1, \dots, b_r) \\ &= 2^r \sum_{n_1, \dots, n_r=0}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} \left(\sum_{j=1}^r a_j n_j + x \right)^k. \end{aligned} \quad (7)$$

By the above Theorem, we may now construct the complex multivariate q -zeta functions as follows:

Definition 1. For $s \in \mathbb{C}$, we define

$$\zeta_r(s, x \mid a_1, \dots, a_r; b_1, \dots, b_r) = \sum_{n_1, \dots, n_r=0}^{\infty} \frac{2^r (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j}}{(\sum_{j=1}^r a_j n_j + x)^s}. \quad (8)$$

Thus we note that this function in (8) is analytic continuation in whole complex plane. And we see this multivariate q -zeta function interpolates q -Euler polynomials at negative integers.

Theorem 3. *Let n be an odd positive integer. Then we have*

$$\zeta_r(-n, x \mid a_1, \dots, a_r; b_1, \dots, b_r) = E_{n,q}^{(r)}(x \mid a_1, \dots, a_r; b_1, \dots, b_r). \quad (9)$$

We now give the complex integral representation of $\zeta_r(s, x \mid a_1, \dots, a_r; b_1, \dots, b_r)$. Using (6), we have the following,

$$\begin{aligned} & \frac{1}{\Gamma(s)} \oint_{\mathbb{C}} F_q^{(r)}(-t, x \mid a_1, \dots, a_r; b_1, \dots, b_r) t^{s-1} dt \\ &= 2^r \sum_{n_1, \dots, n_r=0}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} \frac{1}{\Gamma(s)} \oint_{\mathbb{C}} e^{-(\sum_{j=1}^r a_j n_j + x)t} t^{s-1} dt \\ &= 2^r \sum_{n_1, \dots, n_r=0}^{\infty} (-1)^{\sum_{j=1}^r n_j} \frac{q^{\sum_{j=1}^r b_j n_j}}{(\sum_{j=1}^r a_j n_j + x)^s} \frac{1}{\Gamma(s)} \oint_{\mathbb{C}} e^{-y} y^{s-1} dy \\ &= \zeta_r(s, x \mid a_1, \dots, a_r; b_1, \dots, b_r). \end{aligned} \quad (10)$$

On the other hand

$$\begin{aligned}
& \frac{1}{\Gamma(s)} \oint_{\mathbb{C}} F_q^{(r)}(-t, x \mid a_1, \dots, a_r; b_1, \dots, b_r) t^{s-1} dt \\
&= \sum_{m=0}^{\infty} (-1)^m \frac{E_{m,q}^{(r)}(x \mid a_1, \dots, a_r; b_1, \dots, b_r)}{m!} \frac{1}{\Gamma(s)} \oint_{\mathbb{C}} t^{m+s-1} dt. \quad (11)
\end{aligned}$$

Thus by (10) and (11), we have the following:

$$\begin{aligned}
& \zeta_r(s, x \mid a_1, \dots, a_r; b_1, \dots, b_r) \\
&= \sum_{m=0}^{\infty} (-1)^m \frac{E_m^{(r)}(x \mid a_1, \dots, a_r; b_1, \dots, b_r)}{m!} \frac{1}{\Gamma(s)} \oint_{\mathbb{C}} t^{m+s-1} dt.
\end{aligned}$$

Thus we have

$$\zeta_r(-n, x \mid a_1, \dots, a_r; b_1, \dots, b_r) = E_n^{(r)}(x \mid a_1, \dots, a_r; b_1, \dots, b_r). \quad (12)$$

To construct the multivariate Dirichlet L -function we investigate the generating function of generalized multivariate q -Euler numbers attached to χ , which is derived in (4).

$$\begin{aligned}
& 2^r \sum_{n_1, \dots, n_r=0}^{f-1} \frac{(-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} e^{\sum_{j=1}^r a_j n_j t} \prod_{j=1}^r \chi(n_j)}{\prod_{j=1}^r (q^{f b_j} e^{f a_j t} + 1)} \\
&= 2^r \sum_{n_1, \dots, n_r=0}^{f-1} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} \prod_{j=1}^r \chi(n_j) \\
& \quad \sum_{x_1, \dots, x_r=0}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r f b_j x_j} e^{\sum_{j=1}^r f a_j x_j t} \\
&= 2^r \sum_{x_1, \dots, x_r=0}^{\infty} \sum_{n_1, \dots, n_r=0}^{f-1} (-1)^{\sum_{j=1}^r (n_j + f x_j)} q^{\sum_{j=1}^r b_j (n_j + f x_j)} \\
& \quad \prod_{j=1}^r \chi(n_j + f x_j) e^{t \sum_{j=1}^r a_j (n_j + f x_j)} \\
&= 2^r \sum_{n_1, \dots, n_r=0}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} \prod_{j=1}^r \chi(n_j) e^{t \sum_{j=1}^r a_j n_j}.
\end{aligned}$$

Thus we can write,

$$\begin{aligned}
& F_{\chi,q}^{(r)}(t \mid a_1, \dots, a_r ; b_1, \dots, b_r) \\
&= 2^r \sum_{n_1, \dots, n_r=1}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} \prod_{j=1}^r \chi(n_j) e^{t \sum_{j=1}^r a_j n_j} \\
&\quad (\text{since } \chi(0) = 0, \text{ we can start at } n_1 = 1, n_2 = 1, \dots, n_r = 1) \\
&= \sum_{n=1}^{\infty} E_{n,\chi,q}(a_1, \dots, a_r ; b_1, \dots, b_r) \frac{t^n}{n!}. \tag{13}
\end{aligned}$$

From (13), we can derive the following

$$\begin{aligned}
& 2^r \sum_{n_1, \dots, n_r=1}^{\infty} (-1)^{\sum_{j=1}^r n_j} q^{\sum_{j=1}^r b_j n_j} \prod_{j=1}^r \chi(n_j) (a_1 n_1 + \dots + a_r n_r) \\
&= E_{k,\chi,q}(a_1, \dots, a_r ; b_1, \dots, b_r). \tag{14}
\end{aligned}$$

Therefore we have the following:

Definition 2. For $s \in \mathbb{C}$, define multivariate Dirichlet L -function as follows:

$$\begin{aligned}
& L_r(s, \chi \mid a_1, \dots, a_r ; b_1, \dots, b_r) \\
&= 2^r \sum_{n_1, \dots, n_r=1}^{\infty} \frac{(-1)^{n_1 + \dots + n_r} q^{b_1 n_2 + \dots + b_r n_r} \chi(n_1) \dots \chi(n_r)}{(a_1 n_1 + \dots + a_r n_r)^s}. \tag{15}
\end{aligned}$$

Note that $L_r(s, \chi \mid a_1, \dots, a_r ; b_1, \dots, b_r)$ is also analytic function in whole complex plane. By (12), (13), (14) and (15), we see that the q -analogue multivariate Dirichlet L -function interpolates multivariate generalized q -Euler numbers attached to χ at negative integers as follows;

Theorem 4. Let k be a positive integer. Then we have

$$L_r(-k, \chi \mid a_1, \dots, a_r ; b_1, \dots, b_r) = E_{k,\chi,q}(a_1, \dots, a_r ; b_1, \dots, b_r).$$

References

- [1] L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J., **15**(1948), 987-1000.
- [2] L. Carlitz, *q-Bernoulli and Eulerian numbers*, Trans. Amer. Math. Soc., **76**(1954), 332-350
- [3] M. Cenkci, M.Can, *Some results on q-analogue of the Lerch zeta function*, Adv. Stud. Contemp. Math., **12**(2006), 213-223

- [4] M. Cenkci, M.Can, V. Kurt *p-adic interpolation functions and Kummer-type congruences for q-twisted and q-generalized twisted Euler numbers*, Adv. Stud. Contemp. Math., **9**(2004), 203-216
- [5] H.S. Cho, E. S. Kim, *Translation-invariant p-adic integral on Z_p* , Notes Number Theory Discrete Math., **7**(2001), 70-77
- [6] A. S. Hegazi, M. Mansour, *A note on q-Bernoulli numbers and polynomials*, J. Nonlinear Math. Phys., **13**(2006), 9-18
- [7] T. Kim, *p-adic q-integrals associated with the Changhee-Barnes' q-Bernoulli polynomials*, Integral Transforms Spec. Funct., **15**(2004), 415-420
- [8] T. Kim, S. H. Rim, *Generalized Carlitz's q-Bernoulli numbers in the p-adic number field*, Adv. Stud. Contemp. Math., **2**(2000), 9-19
- [9] T. Kim, *On the analogs of Euler numbers and polynomials associated with p-adic q-integrals on Z_p at $q = -1$* , J. Math. Anal. Appl.(2006) doi:10.1016/j.jmaa.2006.09.027
- [10] T. Kim, *A new approach to q-zeta function*, Adv. Stud. Contemp. Math., **11**(2005), 157-162
- [11] T. Kim, L. C. Jang, H. K. Park, *A note on q-Euler and Genocchi numbers*, Proc. Japan Academy, **77**(2001), 139-141
- [12] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys., **12**(2005), 189-196.
- [13] T. Kim, *Non-Archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials*, Russ. J. Math. Phys., **10**(2003), 91-98.
- [14] T. Kim, *An invariant p-adic integral associated with Daehee numbers*, Integral Transforms and special functions, **13**(2002), 65-69.
- [15] T. Kim, *On a q-analogue of the p-adic log gamma functions and relate integrals*, J. Number Theory, **76**(1999), 320-329.
- [16] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys., **9**(2002), 288-299.
- [17] T. Kim, *q-Euler numbers and polynomials associated with p-adic q-integrals*, J. Nonlinear Math. Phys.(accepted).
- [18] T. Kim, *q-Euler numbers and polynomials associated with p-adic q-integrals and basic q-zeta function*, Trends in Mathematics (Information Center for Mathematical Sciences), **9**(2006), 7-12.
- [19] B. A. Kupershmidt, *Reflection symmetries of q-Bernoulli polynomials*, J. Nonlinear Math. Phys., **12**(2005), 412-422.
- [20] J. Satoh, *q-analogue of Riemann's ζ -function and q-Euler numbers*, J. Number Theory, **31**(1989), 346-362.
- [21] K. Shiratani, S. Yamamoto, *On a p-adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci. Kyushu Univ. Ser. A, **39**(1985), 113-125.
- [22] Y. Simsek, *Theorems on twisted L-function and twisted Bernoulli numbers*, Adv. Stud. Contemp. Math., **11**(2005), 205-218.
- [23] Y. Simsek, D. Kim, S. H. Rim, *On the two-variable Dirichlet q-L-series*, Adv. Stud. Contemp. Math., **10**(2005), 131-142.
- [24] Q.-M. Luo, H. M. Srivastava, *Some relationships between the Apostol-Bernoulli and Apostol-Euler polynomials*, Comput. Math. Appl., **10**(2005), 631-642.